

ALGEBRAIC MONTGOMERY-YANG PROBLEM: THE NONCYCLIC CASE

DONGSEON HWANG AND JONGHAE KEUM

ABSTRACT. Montgomery-Yang problem predicts that every pseudofree circle action on the 5-dimensional sphere has at most 3 non-free orbits. Using a certain one-to-one correspondence, Kollár formulated the algebraic version of the Montgomery-Yang problem: every projective surface S with quotient singularities such that $b_2(S) = 1$ has at most 3 singular points if its smooth locus S^0 is simply-connected.

In this paper, we prove the conjecture under the assumption that S has at least one noncyclic singularity. In the course of the proof, we classify projective surfaces S with quotient singularities such that (i) $b_2(S) = 1$, (ii) $H_1(S^0, \mathbb{Z}) = 0$, and (iii) S has 4 or more singular points, not all cyclic, and prove that all such surfaces have $\pi_1(S^0) \cong \mathfrak{A}_5$, the icosahedral group.

1. INTRODUCTION

A pseudofree $\mathbb{S}^1$ -action on a sphere $\mathbb{S}^{2k-1}$ is a smooth $\mathbb{S}^1$ -action which is free except for finitely many non-free orbits (whose isotropy types $\mathbb{Z}_{m_1}, \dots, \mathbb{Z}_{m_n}$ have pairwise relatively prime orders).

For $k = 2$ Seifert [Se] showed that such an action must be linear and hence has at most two non-free orbits. In the contrast to this, for $k = 4$ Montgomery and Yang [MY] showed that given any pairwise relatively prime collection of positive integers $m_1, \dots, m_n$, there is a pseudofree $\mathbb{S}^1$ -action on homotopy 7-sphere whose non-free orbits have exactly those orders. Petrie [P] proved similar results in all higher odd dimensions. This led Fintushel and Stern to formulate the following problem:

Conjecture 1.1 ([FS87]). (Montgomery-Yang Problem)

Let

$$\mathbb{S}^1 \times \mathbb{S}^5 \rightarrow \mathbb{S}^5$$

be a pseudofree $\mathbb{S}^1$ -action. Then it has at most 3 non-free orbits.

The problem has remained unsolved since its formulation.

Pseudofree $\mathbb{S}^1$ -actions on 5-manifolds L have been studied in terms of the 4-dimensional quotient orbifold $L/\mathbb{S}^1$ (see e.g., [FS85], [FS87]). The following one-to-one correspondence was known to Montgomery, Yang, Fintushel and Stern, and recently observed by Kollár ([Kol05], [Kol08]):

Theorem 1.2. *There is a one-to-one correspondence between:*

2000 *Mathematics Subject Classification.* Primary 14J17.

Key words and phrases. Algebraic Montgomery-Yang Problem, rational homology projective plane, quotient singularity, orbifold Bogomolov-Miyaoka-Yau inequality.

Research supported by Basic Science Research Program through the National Research Foundation(NRF) of Korea funded by the Ministry of education, Science and Technology (2010-0001652).

- (1) Pseudofree $\mathbb{S}^1$ -actions on 5 dimensional rational homology spheres L with $H_1(L, \mathbb{Z}) = 0$.
- (2) Smooth, compact 4-manifolds M with boundary such that
 - (a) $\partial M = \cup_i L_i$ is a disjoint union of lens spaces $L_i = \mathbb{S}^3/\mathbb{Z}_{m_i}$,
 - (b) the m_i are relatively prime to each other,
 - (c) $H_1(M, \mathbb{Z}) = 0$ and $H_2(M, \mathbb{Z}) \cong \mathbb{Z}$.

Furthermore, L is diffeomorphic to $\mathbb{S}^5$ iff $\pi_1(M) = 1$.

Using the one-to-one correspondence, Kollár formulated the algebraic version of the Montgomery-Yang problem as follows:

Conjecture 1.3. [Kol08] (Algebraic Montgomery-Yang Problem)

Let S be a rational homology projective plane with quotient singularities, i.e., a normal projective surface with quotient singularities such that $b_2(S) = 1$. Assume that $S^0 := S \setminus \text{Sing}(S)$ is simply-connected. Then S has at most 3 singular points.

In this paper, we verify the conjecture when S has at least one noncyclic singularity. More precisely, we prove the following:

Theorem 1.4. Let S be a rational homology projective plane with quotient singularities such that $\pi_1(S^0) = \{1\}$. Assume that S has at least one noncyclic singularity. Then $|\text{Sing}(S)| \leq 3$.

We note that the condition $\pi_1(S^0) = \{1\}$ cannot be replaced by the weaker condition $H_1(S^0, \mathbb{Z}) = 0$. There are infinitely many examples of rational homology projective planes with exactly 4 quotient singularities, 3 cyclic, 1 noncyclic, such that $H_1(S^0, \mathbb{Z}) = 0$ but $\pi_1(S^0) \neq \{1\}$ ([Br] or [Kol08], Example 31). These examples are the global quotients

$$S_{I_m} := \mathbb{CP}^2/I_m = (\mathbb{CP}^2/Z)/\mathfrak{A}_5,$$

where $I_m \subset GL(2, \mathbb{C})$ is the group of order $120m$ in Brieskorn's list (see Table 1), an extension of the icosahedral group $\mathfrak{A}_5 \subset PGL(2, \mathbb{C})$ by the cyclic group $Z \cong \mathbb{Z}_{2m}$, and the action of I_m on $\mathbb{CP}^2$ is induced from the natural action on $\mathbb{C}^2$. We call S_{I_m} a *Brieskorn quotient*.

On the other hand, it follows from the orbifold Bogomolov-Miyaoka-Yau inequality that every rational homology projective plane S with quotient singularities such that $H_1(S^0, \mathbb{Z}) = 0$ has at most 4 singular points (cf. [Kol08], [HK], [Keu10]). Therefore, to prove Theorem 1.4, it is enough to classify rational homology projective planes S with 4 quotient singularities, not all cyclic, such that $H_1(S^0, \mathbb{Z}) = 0$. It turns out that such a surface is deformation equivalent to a Brieskorn quotient.

Theorem 1.5. Let S be a rational homology projective plane with 4 quotient singularities, not all cyclic, such that $H_1(S^0, \mathbb{Z}) = 0$. Then the following hold true.

- (1) S has 3 cyclic singularities of type $\mathbb{C}^2/\mathbb{Z}_2$, $\mathbb{C}^2/\mathbb{Z}_3$, $\mathbb{C}^2/\mathbb{Z}_5$, and one noncyclic singularity of type $\mathbb{C}^2/I_m$, where $I_m \subset GL(2, \mathbb{C})$ is the $2m$ -ary icosahedral group of order $120m$ (in Brieskorn's notation). Furthermore, the 3 cyclic singularities are of type $\frac{1}{2}(1, 1)$, $\frac{1}{3}(1, \alpha)$, $\frac{1}{5}(1, \beta)$, if the 3 branches of the dual graph of the noncyclic singularity are of type $\frac{1}{2}(1, 1)$, $\frac{1}{3}(1, 3 - \alpha)$, $\frac{1}{5}(1, 5 - \beta)$ (see Table 4).
- (2) $-K_S$ is ample.

- (3) *The minimal resolution of S can be obtained by starting with a minimal rational ruled surface and blowing up inside 3 of the fibres, i.e. the blowing up starts at three centers, one on each of the 3 fibres.*
- (4) *S^0 is deformation equivalent to $(\mathbb{CP}^2/I_m)^0$, where I_m is determined by the noncyclic singularity of S and its action on $\mathbb{CP}^2$ is induced by the natural action on $\mathbb{C}^2$. The deformation space has dimension 2.*
- (5) *$\pi_1(S^0) \cong \mathfrak{A}_5$, the alternating group of order 60.*

We will prove Theorem 1.5. In the proof, we use the orbifold Bogomolov-Miyaoka-Yau inequality (Theorem 2.2 and 2.3) and a detailed computation for (-1) -curves on the minimal resolution S' of S . The latter idea was used in [Kou08].

Remark 1.6. Consider a Brieskorn quotient $S_{I_m} := \mathbb{CP}^2/I_m = (\mathbb{CP}^2/Z)/\mathfrak{A}_5$. The cone $\mathbb{CP}^2/Z$ is the closure of the $\mathfrak{A}_5$ -universal cover of $S_{I_m}^0$. Note that the cone has no deformation. Thus the deformation of $S_{I_m}^0$ must correspond to a deformation of the I_m -action on $\mathbb{CP}^2$. This was pointed out to us by János Kollár. It is an interesting problem to describe explicitly such a deformation.

Throughout this paper, we work over the field $\mathbb{C}$ of complex numbers.

2. ALGEBRAIC SURFACES WITH QUOTIENT SINGULARITIES

2.1. A singularity p of a normal surface S is called a quotient singularity if locally the germ is analytically isomorphic to $(\mathbb{C}^2/G, O)$ for some nontrivial finite subgroup G of $GL_2(\mathbb{C})$ without quasi-reflections. Brieskorn classified such finite subgroups of $GL(2, \mathbb{C})$ [Bri]. Table 1 summarizes the result. Here we only explain the notation for dual graph.

$$\begin{aligned}
 \langle q, q_1 \rangle &:= \text{the dual graph of the singularity of type } \frac{1}{q}(1, q_1) \\
 \langle b; s_1, t_1; s_2, t_2; s_3, t_3 \rangle &:= \text{the tree of the form} \\
 &\quad \begin{array}{c} \langle s_2, t_2 \rangle \\ \downarrow \\ \langle s_1, t_1 \rangle - \underset{-b}{\circ} - \langle s_3, t_3 \rangle \end{array}
 \end{aligned}$$

For more information about the table, we refer to the original paper of Brieskorn[Bri].

2.2. Let S be a normal projective surface with quotient singularities and

$$f : S' \rightarrow S$$

be a minimal resolution of S . It is well-known that quotient singularities are log-terminal singularities. Thus one can write

$$K_{S'} \equiv_{\text{num}} f^* K_S - \sum_{p \in \text{Sing}(S)} \mathcal{D}_p$$

where, for each singular point p , $\mathcal{D}_p = \sum (a_j E_j)$ is an effective $\mathbb{Q}$ -divisor supported on $f^{-1}(p) = \cup E_j$ with $0 \leq a_j < 1$. It implies that

$$K_S^2 = K_{S'}^2 - \sum_p \mathcal{D}_p^2 = K_{S'}^2 + \sum_p \mathcal{D}_p K_{S'}.$$

TABLE 1. Classification of finite subgroups of $GL(2, \mathbb{C})$

Type	G	$ G $	$G/[G, G]$	Dual Graph Γ_G
A_{q, q_1}	C_{q, q_1}	q	$\mathbb{Z}_q$	$\langle q, q_1 \rangle$ $0 < q_1 < q, (q, q_1) = 1$
D_{q, q_1}	$(Z_{2m}, Z_{2m}; D_q, D_q)$	$4mq$	$\mathbb{Z}_{2m} \times \mathbb{Z}_2$	$\langle b; 2, 1; 2, 1; q, q_1 \rangle$ $m = (b-1)q - q_1$ odd
D_{q, q_1}	$(Z_{4m}, Z_{2m}; D_q, C_{2q})$	$4mq$	$\mathbb{Z}_{4m}$	$\langle b; 2, 1; 2, 1; q, q_1 \rangle$ $m = (b-1)q - q_1$ even
T_m	$(Z_{2m}, Z_{2m}; T, T)$	$24m$	$\mathbb{Z}_{3m}$	$\langle b; 2, 1; 3, 2; 3, 2 \rangle, m = 6(b-2) + 1$ $\langle b; 2, 1; 3, 1; 3, 1 \rangle, m = 6(b-2) + 5$
T_m	$(Z_{2m}, Z_{2m}; T, D_2)$	$24m$	$\mathbb{Z}_{3m}$	$\langle b; 2, 1; 3, 1; 3, 2 \rangle, m = 6(b-2) + 3$
O_m	$(Z_{2m}, Z_{2m}; O, O)$	$48m$	$\mathbb{Z}_{2m}$	$\langle b; 2, 1; 3, 2; 4, 3 \rangle, m = 12(b-2) + 1$ $\langle b; 2, 1; 3, 1; 4, 3 \rangle, m = 12(b-2) + 5$ $\langle b; 2, 1; 3, 2; 4, 1 \rangle, m = 12(b-2) + 7$ $\langle b; 2, 1; 3, 1; 4, 1 \rangle, m = 12(b-2) + 11$
I_m	$(Z_{2m}, Z_{2m}; I, I)$	$120m$	$\mathbb{Z}_m$	$\langle b; 2, 1; 3, 2; 5, 4 \rangle, m = 30(b-2) + 1$ $\langle b; 2, 1; 3, 2; 5, 3 \rangle, m = 30(b-2) + 7$ $\langle b; 2, 1; 3, 1; 5, 4 \rangle, m = 30(b-2) + 11$ $\langle b; 2, 1; 3, 2; 5, 2 \rangle, m = 30(b-2) + 13$ $\langle b; 2, 1; 3, 1; 5, 3 \rangle, m = 30(b-2) + 17$ $\langle b; 2, 1; 3, 2; 5, 1 \rangle, m = 30(b-2) + 19$ $\langle b; 2, 1; 3, 1; 5, 2 \rangle, m = 30(b-2) + 23$ $\langle b; 2, 1; 3, 1; 5, 1 \rangle, m = 30(b-2) + 29$

Lemma 2.1. *If $-K_S$ is ample, then $C^2 \geq -1$ for any irreducible curve $C \subset S'$ not contracted by $f : S' \rightarrow S$.*

Proof. Note that $C(f^*K_S) < 0$ and $C(\sum \mathcal{D}_p) \geq 0$. Thus $CK_{S'} < 0$, and hence $C^2 \geq -1$. $\square$

Also we recall the orbifold Euler characteristic

$$e_{orb}(S) := e(S) - \sum_{p \in \text{Sing}(S)} \left(1 - \frac{1}{|G_p|}\right)$$

where G_p is the local fundamental group of p .

The following theorem, called the orbifold Bogomolov-Miyaoka-Yau inequality, is one of the main ingredients in the proof of our main theorem.

Theorem 2.2 ([S], [Mi], [KNS], [Me]). *Let S be a normal projective surface with quotient singularities such that K_S is nef. Then*

$$K_S^2 \leq 3e_{orb}(S).$$

We also need the following weaker inequality, which also holds when K_S is nef.

Theorem 2.3 ([KM]). *Let S be a normal projective surface with quotient singularities such that $-K_S$ is nef. Then*

$$0 \leq e_{orb}(S).$$

2.3. Let S be a normal projective surface with quotient singularities and $f : S' \rightarrow S$ be a minimal resolution of S . It is well-known that the torsion-free part of the second cohomology group,

$$H^2(S', \mathbb{Z})_{free} := H^2(S', \mathbb{Z})/torsion$$

has a lattice structure which is unimodular. For a quotient singular point $p \in S$, let

$$R_p \subset H^2(S', \mathbb{Z})_{free}$$

be the sublattice of $H^2(S', \mathbb{Z})_{free}$ spanned by the numerical classes of the components of $f^{-1}(p)$. It is a negative definite lattice, and its discriminant group

$$\text{disc}(R_p) := \text{Hom}(R_p, \mathbb{Z})/R_p$$

is isomorphic to the abelianization $G_p/[G_p, G_p]$ of the local fundamental group G_p . In particular, the absolute value $|\det(R_p)|$ of the determinant of the intersection matrix of R_p is equal to the order $|G_p/[G_p, G_p]|$. Let

$$R = \bigoplus_{p \in \text{Sing}(S)} R_p \subset H^2(S', \mathbb{Z})_{free}$$

be the sublattice of $H^2(S', \mathbb{Z})_{free}$ spanned by the numerical classes of the exceptional curves of $f : S' \rightarrow S$. We also consider the sublattice

$$R + \langle K_{S'} \rangle \subset H^2(S', \mathbb{Z})_{free}$$

spanned by R and the canonical class $K_{S'}$. Note that

$$\text{rank}(R) \leq \text{rank}(R + \langle K_{S'} \rangle) \leq \text{rank}(R) + 1.$$

Lemma 2.4 ([HK], Lemma 3.3). *Let S be a normal projective surface with quotient singularities and $f : S' \rightarrow S$ be a minimal resolution of S . Then the following hold true.*

- (1) $\text{rank}(R + \langle K_{S'} \rangle) = \text{rank}(R)$ if and only if K_S is numerically trivial.
- (2) $\det(R + \langle K_{S'} \rangle) = \det(R) \cdot K_S^2$ if K_S is not numerically trivial.
- (3) If in addition $b_2(S) = 1$ and K_S is not numerically trivial, then $R + \langle K_{S'} \rangle$ is a sublattice of finite index in the unimodular lattice $H^2(S', \mathbb{Z})_{free}$, in particular $|\det(R + \langle K_{S'} \rangle)|$ is a nonzero square number.

We denote the number $|\det(R + \langle K_{S'} \rangle)|$ by D , i.e., we define

$$D := |\det(R + \langle K_{S'} \rangle)|.$$

The following will be also used in our proof.

Lemma 2.5. *Let S be a rational homology projective plane with quotient singularities such that $H_1(S^0, \mathbb{Z}) = 0$. Let $f : S' \rightarrow S$ be a minimal resolution. Then*

- (1) $H^2(S', \mathbb{Z})$ is torsion free, i.e. $H^2(S', \mathbb{Z}) = H^2(S', \mathbb{Z})_{free}$,
- (2) R is a primitive sublattice of the unimodular lattice $H^2(S', \mathbb{Z})$,
- (3) $\text{disc}(R)$ is a cyclic group, in particular, the orders $|G_p/[G_p, G_p]| = |\det(R_p)|$ are pairwise relatively prime,
- (4) K_S is not numerically trivial, i.e. K_S is either ample or anti-ample,
- (5) $D = |\det(R)|K_S^2$ and D is a nonzero square number,

- (6) the Picard group $Pic(S')$ is generated over $\mathbb{Z}$ by the exceptional curves and a $\mathbb{Q}$ -divisor M of the form

$$M = \frac{1}{\sqrt{D}} f^* K_S + z,$$

where z is a generator of $\text{disc}(R)$, hence of the form $z = \sum_{p \in \text{Sing}(S)} b_p e_p$ for some integers b_p , where e_p is a generator of $\text{disc}(R_p)$.

Proof. (1), (2) and (3) are easy to see. cf. [Kou07], Proposition 2.3 and Lemma 3.4.

(4) Assume that K_S is numerically trivial. Then S' is an Enriques surface if all singularities are rational double points, and is a rational surface otherwise. If S' is an Enriques surface, then $H_1(S', \mathbb{Z}) \neq 0$ since $H_1(S', \mathbb{Z}) = \mathbb{Z}/2$ (cf. Proposition 2.3 in [Kou07]). Thus S is a rational surface, and

$$K_{S'} \equiv_{\text{num}} - \sum_{p \in \text{Sing}(S)} \mathcal{D}_p$$

with $\mathcal{D}_p \not\equiv_{\text{num}} 0$ for some p . Note that $\mathcal{D}_p$ defines an element of $R_p^* := \text{Hom}(R_p, \mathbb{Z})$ and the discriminant group $\text{disc}(R_p) := R_p^*/R_p$ has order $|\det(R_p)|$. Thus $|\det(R_p)|\mathcal{D}_p \in R_p$ but $\mathcal{D}_p \notin R_p$ if $\mathcal{D}_p \not\equiv_{\text{num}} 0$. Now we see that

$$\left(\prod_p |\det(R_p)| \right) K_{S'} \in R \subset H^2(S', \mathbb{Z})$$

but $K_{S'} \notin R$. Hence the primitive closure $\bar{R}$ of R in $H^2(S', \mathbb{Z})$ is not equal to R . Now by Lemma 2.5 in [Kou07], $H_1(S^0, \mathbb{Z}) \neq 0$.

(5) follows from (4) and Lemma 2.4.

(6) Note first that $Pic(S') = H^2(S', \mathbb{Z})$ and the sublattice $R \subset H^2(S', \mathbb{Z})$ generated by the exceptional curves is a primitive sublattice of corank 1. Let $R^\perp \subset H^2(S', \mathbb{Z})$ be the orthogonal complement of R . Note that $R^\perp$ is positive definite and of rank 1. Since $H^2(S', \mathbb{Z})$ is unimodular,

$$\det(R^\perp) = |\det(R)| = \prod_{p \in \text{Sing}(S)} |\det(R_p)|.$$

Note that $f^* K_S \in R^\perp$. Thus $R^\perp$ is generated by

$$v = \frac{|\det(R)|}{\sqrt{D}} f^* K_S,$$

and $\text{disc}(R^\perp)$ is generated by $\frac{1}{\sqrt{D}} f^* K_S$. Also note that

$$\text{disc}(R^\perp \oplus R) \cong (\mathbb{Z}/|\det(R)|) \oplus (\mathbb{Z}/|\det(R)|).$$

Thus $Pic(S')/(R^\perp \oplus R)$ is an isotropic subgroup of $\text{disc}(R^\perp \oplus R)$ of order $|\det(R)|$, hence is generated by an element $M \in \text{disc}(R^\perp \oplus R)$ of order $|\det(R)|$. Moreover M is the sum of a generator of $\text{disc}(R^\perp)$ and a generator of $\text{disc}(R)$, since $Pic(S')$ is unimodular. By replacing M by kM for a suitable choice of an integer k , we get M of the desired form

$$M = \frac{1}{\sqrt{D}} f^* K_S + \sum_{p \in \text{Sing}(S)} a_p e_p$$

for some integers a_p , $0 \leq a_p < |\det(R_p)|$. This implies that $\text{Pic}(S')$ is generated over $\mathbb{Z}$ by R , v and M . Finally, note

$$|\det(R)|M = v \pmod{R},$$

a generator of $R^\perp$. Thus $\text{Pic}(S')$ is generated over $\mathbb{Z}$ by R and M . $\square$

3. PROOF OF THEOREM 1.5

Let S be a rational homology projective plane with 4 or more quotient singularities with $H_1(S^0, \mathbb{Z}) = 0$. By Lemma 2.5(3), the orders of the abelianized local fundamental groups are pairwise relatively prime. Thus by Theorem 2.3, one can see that S has 4 singular points and the 4-tuple of orders of the local fundamental groups must be one of the following:

- (1) $(2, 3, 5, q)$, $q \geq 7$,
- (2) $(2, 3, 7, q)$, $11 \leq q \leq 41$,
- (3) $(2, 3, 11, 13)$.

Table 1 shows that all noncyclic singularities of type different from I_m have abelianized local fundamental groups of order divisible by 2 or 3.

Assume that one of the singularities is noncyclic. By Lemma 2.5(3), it must be of type I_m and the other 3 singularities are cyclic of order 2, 3 and 5, respectively. Here we recall that $I_m \subset GL(2, \mathbb{C})$ is the $2m$ -ary icosahedral group of order $120m$. Table 1 shows that there are 8 infinite cases of type I_m .

There are two types of order 3, $< 3, 2 >$ and $< 3, 1 >$; three types of order 5, $< 5, 4 >$, $< 5, 3 > \cong < 5, 2 >$ and $< 5, 1 >$. Thus there are exactly 48 infinite cases for possible combinations of types of singularities. That is, there are exactly 48 infinite cases for R , the sublattice of $\text{Pic}(S') = H^2(S', \mathbb{Z})$ generated by all exceptional curves, where $f : S' \rightarrow S$ is a minimal resolution. In each of the 48 cases we compute $D = |\det(R)|K_S^2$ and check if D is a square number (see Lemma 2.5(5)), using elementary number theoretic arguments. There remain 8 infinite cases and 2 sporadic cases, as given in Table 2 and Table 3. In both tables, the entries of the column b are the possible values of b that make D a square number.

We will explain how to compute D . First note that

$$|\det(R)| = 2 \cdot 3 \cdot 5 \cdot m = 30m.$$

To compute K_S^2 , we use the equality from (2.2)

$$K_S^2 = K_{S'}^2 + \sum_p \mathcal{D}_p K_{S'}.$$

Note that S' has $H^1(S', \mathcal{O}_{S'}) = H^2(S', \mathcal{O}_{S'}) = 0$. Thus by Noether formula,

$$K_{S'}^2 = 12 - e(S') = 10 - b_2(S') = 9 - \mu$$

where μ is the number of the exceptional curves of f .

For each singular point p , the coefficients of the $\mathbb{Q}$ -divisor $\mathcal{D}_p$ can be obtained by solving the equations given by the adjunction formula

$$\mathcal{D}_p E = -K_{S'} E = 2 + E^2$$

for each exceptional curve $E \subset f^{-1}(p)$. Once we know the coefficients, we can easily compute the intersection number $\mathcal{D}_p K_{S'}$.

We first rule out the two sporadic cases.

TABLE 2.

Type of R	$D = \det(R) K_S^2$	b
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 2; 5, 4 \rangle$	$180(5b^2 - 50b + 79)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 2; 5, 3 \rangle$	$180(5b^2 - 36b + 48)$	$b = 8$
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 1; 5, 4 \rangle$	$180(5b^2 - 40b + 52)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 2; 5, 2 \rangle$	$180(5b^2 - 34b + 41)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 1; 5, 3 \rangle$	$180(5b^2 - 26b + 27)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 2; 5, 1 \rangle$	$180(5b^2 - 20b + 18)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 1; 5, 2 \rangle$	$180(5b^2 - 24b + 22)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 1; 5, 1 \rangle$	$900(b - 1)^2$	$b \geq 2$
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 2; 5, 4 \rangle$	$36(25b^2 - 190b + 277)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 2; 5, 3 \rangle$	$36(25b^2 - 120b + 134)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 1; 5, 4 \rangle$	$36(25b^2 - 140b + 162)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 2; 5, 2 \rangle$	$36(25b^2 - 110b + 111)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 1; 5, 3 \rangle$	$36(5b - 7)^2$	$b \geq 2$
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 2; 5, 1 \rangle$	$36(25b^2 - 40b + 8)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 1; 5, 2 \rangle$	$36(5b - 6)^2$	$b \geq 2$
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 1; 5, 1 \rangle$	$36(25b^2 + 10b - 37)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 2; 5, 4 \rangle$	$36(25b^2 - 130b + 159)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 2; 5, 3 \rangle$	$36(25b^2 - 60b + 28)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 1; 5, 4 \rangle$	$36(5b - 8)^2$	$b \geq 2$
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 2; 5, 2 \rangle$	$36(25b^2 - 50b + 17)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 1; 5, 3 \rangle$	$36(25b^2 - 10b - 37)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 2; 5, 1 \rangle$	$36(25b^2 + 20b - 74)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 1; 5, 2 \rangle$	$36(25b^2 - 38)$	none
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 1; 5, 1 \rangle$	$36(25b^2 + 70b - 99)$	none

Lemma 3.1. *The case $\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 4 \rangle + \langle 8; 2, 1; 3, 2; 5, 3 \rangle$ does not occur.*

Proof. In this case, $m = 30(b - 2) + 7 = 187$, so

$$|\det(R)| = 30 \cdot 187.$$

The number of exceptional curves $\mu = 13$, so $K_{S'}^2 = -4$, where $f : S' \rightarrow S$ is a minimal resolution. Let p_1, p_2, p_3, p_4 be the four singular points. Let $E_1, \dots, E_6$ be

TABLE 3.

Type of R	$D = \det(R) K_S^2$	b
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 2; 5, 4 \rangle$	$20(45b^2 - 390b + 593)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 2; 5, 3 \rangle$	$20(45b^2 - 264b + 326)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 1; 5, 4 \rangle$	$100(9b^2 - 60b + 74)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 2; 5, 2 \rangle$	$20(45b^2 - 246b + 275)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 1; 5, 3 \rangle$	$20(45b^2 - 174b + 157)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 2; 5, 1 \rangle$	$100(3b - 4)^2$	$b \geq 2$
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 1; 5, 2 \rangle$	$20(45b^2 - 156b + 124)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 1; 5, 1 \rangle$	$20(45b^2 - 30b - 17)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 2; 5, 4 \rangle$	$4(225b^2 - 1410b + 1903)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 2; 5, 3 \rangle$	$4(15b - 26)^2$	$b \geq 2$
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 1; 5, 4 \rangle$	$4(225b^2 - 960b + 968)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 2; 5, 2 \rangle$	$4(15b - 23)^2$	$b \geq 2$
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 1; 5, 3 \rangle$	$4(225b^2 - 330b + 11)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 2; 5, 1 \rangle$	$4(225b^2 - 60b - 338)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 1; 5, 2 \rangle$	$4(225b^2 - 240b - 46)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 1; 5, 1 \rangle$	$4(225b^2 + 390b - 643)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 2; 5, 4 \rangle$	$4(15b - 29)^2$	$b \geq 2$
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 2; 5, 3 \rangle$	$4(225b^2 - 240b - 278)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 1; 5, 4 \rangle$	$4(225b^2 - 420b + 86)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 2; 5, 2 \rangle$	$4(225b^2 - 150b - 317)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 1; 5, 3 \rangle$	$4(225b^2 + 210b - 763)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 2; 5, 1 \rangle$	$4(225b^2 + 480b - 1076)$	$b = 2$
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 1; 5, 2 \rangle$	$4(225b^2 + 300b - 712)$	none
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 1; 5, 1 \rangle$	$4(225b^2 + 930b - 1201)$	none

the components of $f^{-1}(p_4)$ such that

$$\begin{array}{c}
-2 \\
E_2 - E_3 - E_6 - E_5 - E_4 \\
\downarrow \\
E_1 \\
-2
\end{array}$$

Solving the equations given by the adjunction formula, we get

$$K_{S'} = f^*K_S - \frac{93E_1 + 186E_6 + 62E_2 + 124E_3 + 112E_4 + 149E_5}{187}.$$

It is easy to compute that

$$K_S^2 = K_{S'}^2 + \frac{186E_6K_{S'} + 112E_4K_{S'}}{187} = -4 + \frac{186 \cdot 6 + 112}{187} = \frac{480}{187}.$$

Thus

$$D = |\det(R)|K_S^2 = 120^2.$$

Note that $K_S^2 > 3e_{orb}(S)$, so $-K_S$ is ample by the orbifold Bogomolov-Miyaoka-Yau inequality. Thus S' is a rational surface, not minimal. Also note that the divisor M from Lemma 2.5(6) takes the form

$$M = \frac{1}{120}f^*K_S + \sum_{p \in \text{Sing}(S)} a_p e_p.$$

Let C be a (-1) -curve on S' . By Lemma 2.5(6), C can be written as

$$C = kM + r$$

for some integer k and some $r \in R$, hence as

$$C = \frac{k}{120}f^*K_S + C(1) + C(2) + C(3) + C(4)$$

where $C(i)$ is a $\mathbb{Q}$ -divisor supported on $f^{-1}(p_i)$. Note that

$$C^2 = \left(\frac{k}{120}f^*K_S\right)^2 + C(1)^2 + C(2)^2 + C(3)^2 + C(4)^2.$$

Since $(f^*K_S)C(i) = 0$ for all i , we have

$$(f^*K_S)C = (f^*K_S)\left(\frac{k}{120}f^*K_S\right) = \frac{k}{120}K_S^2 = \frac{4k}{187}.$$

Since $-K_S$ is ample and $C \notin R$, we see that $(f^*K_S)C < 0$, hence

$$k < 0.$$

Note that $K_{S'}C = -1$. From the equality

$$K_{S'}C = (f^*K_S)C - \frac{(93E_1 + 186E_6 + 62E_2 + 124E_3 + 112E_4 + 149E_5)C}{187},$$

we get

$$(93E_1 + 186E_6 + 62E_2 + 124E_3 + 112E_4 + 149E_5)C = 187 + 4k.$$

This is possible only if

$$E_6C = E_5C = E_4C = E_3C = 0, \quad E_2C = E_1C = 1, \quad k = -8.$$

Since $E_jC(4) = E_jC$ for $j = 1, \dots, 6$, we obtain the coefficients of $C(4)$ by solving the equations given by the above intersection numbers.

$$C(4) = -\frac{106E_1 + 133E_2 + 79E_3 + 5E_4 + 15E_5 + 25E_6}{187} = E_1^* + E_2^*,$$

where $E_j^* \in \text{Hom}(R_{p_4}, \mathbb{Z})$ is the dual vector of E_j . Thus

$$C(4)^2 = (E_1^* + E_2^*)C(4) = -\frac{106 + 133}{187}.$$

Now we have

$$\sum_{j \leq 3} C(j)^2 = C^2 - C(4)^2 - \left(\frac{-8f^*K_S}{120}\right)^2 = -1 + \frac{239}{187} - \frac{32}{15 \cdot 187} > 0$$

which contradicts the negative definiteness of exceptional curves. $\square$

Lemma 3.2. *The case $\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 1 \rangle + \langle 2; 2, 1; 3, 2; 5, 1 \rangle$ does not occur.*

Proof. The proof is similar to the previous case. In this case, $m = 19$ and $\mu = 8$, so $|\det(R)| = 30 \cdot 19$ and $K_S^2 = 1$. Let B_2, B_3 be the components of $f^{-1}(p_2), f^{-1}(p_3)$, $E_1, \dots, E_5$ be the components of $f^{-1}(p_4)$ such that

$$\begin{array}{c} \begin{array}{c} -2 \\ E_2 \end{array} - \begin{array}{c} -2 \\ E_3 \end{array} - \begin{array}{c} -2 \\ E_5 \end{array} - \begin{array}{c} -5 \\ E_4 \end{array} \\ \downarrow \\ E_1 \\ -2 \end{array}$$

Then

$$K_{S'} = f^*K_S - \frac{B_2}{3} - \frac{3B_3}{5} - \frac{9E_1 + 6E_2 + 12E_3 + 15E_4 + 18E_5}{19},$$

$$K_S^2 = \frac{28 \cdot 56}{15 \cdot 19}, \quad D = |\det(R)|K_S^2 = 56^2.$$

Here again by the orbifold Bogomolov-Miyaoka-Yau inequality, $-K_S$ is ample and S' is a rational surface, not minimal. Let C be a (-1) -curve on S' . Then

$$C = \frac{k}{56} f^*K_S + C(1) + C(2) + C(3) + C(4)$$

for some integer k and some $\mathbb{Q}$ -divisor $C(i)$ supported on $f^{-1}(p_i)$.

Since $(f^*K_S)C = \frac{28k}{285} < 0$, we see that $k < 0$ and we get

$$95B_2C + 171B_3C + 15(9E_1 + 6E_2 + 12E_3 + 15E_4 + 18E_5)C = 285 + 28k.$$

This is impossible because $k < 0$ and $E_jC \geq 0, B_iC \geq 0$ for every i, j . $\square$

Lemma 3.3. *For any of the 8 infinite cases, $-K_S$ is ample.*

Proof. For the 8 infinite cases, we compute K_S^2 as follows.

TABLE 4.

Type of R	K_S^2
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 1; 5, 1 \rangle$	$\frac{30(b-1)^2}{30b-31} \geq \frac{30}{29}$
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 2 \rangle + \langle b; 2, 1; 3, 1; 5, 3 \rangle$	$\frac{6(5b-7)^2}{5(30b-43)} \geq \frac{54}{85}$
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 1; 5, 2 \rangle$	$\frac{6(5b-6)^2}{5(30b-37)} \geq \frac{96}{115}$
$\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 1; 5, 4 \rangle$	$\frac{6(5b-8)^2}{5(30b-49)} \geq \frac{24}{55}$
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 2; 5, 1 \rangle$	$\frac{10(3b-4)^2}{3(30b-41)} \geq \frac{40}{57}$
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 2 \rangle + \langle b; 2, 1; 3, 2; 5, 3 \rangle$	$\frac{2(15b-26)^2}{15(30b-53)} \geq \frac{32}{105}$
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 2; 5, 2 \rangle$	$\frac{2(15b-23)^2}{15(30b-47)} \geq \frac{98}{195}$
$\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 2; 5, 4 \rangle$	$\frac{2(15b-29)^2}{15(30b-59)} \geq \frac{2}{15}$

In each case, $e_{orb}(S) = -1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{5} + \frac{1}{120m} \leq \frac{5}{120}$. From the table we see that $K_S^2 > 3e_{orb}(S)$, so $-K_S$ is ample by the orbifold Bogomolov-Miyaoka-Yau inequality. $\square$

This completes the proof of (1) and (2) of Theorem 1.5. To prove the remaining part, we need to analyze (-1) -curves on the minimal resolution S' . Note that by Lemma 2.1 S' contains no $(-n)$ -curve with $n \geq 2$ other than the exceptional curves of $f : S' \rightarrow S$.

The following proposition will be proved case by case in the next section.

Proposition 3.4. *If S has 4 singularities p_1, p_2, p_3, p_4 of type $\langle 2, 1 \rangle, \langle 3, \alpha \rangle, \langle 5, \beta \rangle, \langle b; 2, 1; 3, 3 - \alpha; 5, 5 - \beta \rangle$, $b \geq 2$, respectively, as in Table 4, then there are three mutually disjoint (-1) -curves C_1, C_2, C_3 on S' such that*

- (1) *each C_i intersects exactly 2 components of $f^{-1}(p_1) \cup f^{-1}(p_2) \cup f^{-1}(p_3) \cup f^{-1}(p_4)$ with multiplicity 1 each,*
- (2) *C_1 intersects the component of the branch $\langle 2, 1 \rangle$ of $f^{-1}(p_4)$ and the component of $f^{-1}(p_1)$, C_2 intersects the terminal component of the branch $\langle 3, 3 - \alpha \rangle$ of $f^{-1}(p_4)$ and one end component of $f^{-1}(p_2)$, and C_3 intersects the terminal component of the branch $\langle 5, 5 - \beta \rangle$ of $f^{-1}(p_4)$ and one end component of $f^{-1}(p_3)$ which is a (-2) -curve if $\beta = 2$ or 4, a (-3) -curve if $\beta = 3$, and a (-5) -curve if $\beta = 1$.*

Proposition 3.5. (1) *The surface S' can be blown down to the Hirzebruch surface F_b . Conversely, S' can be obtained by starting with F_b and blowing up inside 3 of the fibres, i.e. the blowing up starts at three centers, one on each of the 3 fibres.*

- (2) *If two rational homology projective planes S_1 and S_2 have the same type of singularities $\langle 2, 1 \rangle + \langle 3, \alpha \rangle + \langle 5, \beta \rangle + \langle b; 2, 1; 3, 3 - \alpha; 5, 5 - \beta \rangle$, $b \geq 2$, then S_1^0 and S_2^0 are deformation equivalent.*

Proof. (1) By Proposition 3.4 there are three mutually disjoint (-1) -curves C_1, C_2, C_3 on S' satisfying (1) and (2) of Proposition 3.4. By starting with them, we can blow down S' to F_b . Furthermore, the blow up process from F_b to S' is carried out inside 3 of the fibres of F_b .

(2) The blow up process from F_b to S' depends on the choice of three fibres, each with a point marked. The three marked points are the centers of the blowing up. The choice of three fibres is unique up to automorphisms of F_b , while the choice of three points, one on each of the fixed three fibres, is not unique up to automorphisms of F_b , but depends on a 2-dimensional moduli. $\square$

This completes the proof of (3) of Theorem 1.5.

The following examples mentioned in Introduction were discussed in [Kol08], Example 31.

Example 3.6. Consider the $2m$ -ary icosahedral group

$$I_m \subset GL(2, \mathbb{C})$$

of order $120m$ in Brieskorn's list (Table 1). Let $Z \subset I_m$ be its center, then $Z \cong \mathbb{Z}_{2m}$ and $I_m/Z \cong \mathfrak{A}_5 \subset PGL(2, \mathbb{C})$, the icosahedral group. Extend the natural I_m -action on $\mathbb{C}^2$ to $\mathbb{CP}^2$. The center acts trivially on the line at infinity and $\mathbb{CP}^2/Z$ is a cone over the rational normal curve of degree $2m = |Z|$. Then

$$S_{I_m} := \mathbb{CP}^2/I_m = (\mathbb{CP}^2/Z)/\mathfrak{A}_5$$

has 4 quotient singularities, one of type $\mathbb{C}^2/I_m$ at the origin, three of order 2, 3, 5 at infinity. The fundamental group of $S_{I_m}^0$ is $\mathfrak{A}_5$. By Theorem 1.5 (1), the types

of the 3 cyclic singularities are determined by the types of the 3 branches of the non-cyclic singularity. By Proposition 3.4 and 3.5, its minimal resolution S'_{I_m} can be blown down to the Hirzebruch surface F_b . Conversely, S'_{I_m} can be obtained by starting with F_b and blowing up inside 3 of the fibres. Here the 3 centers of the blowing up lie on a section of F_b .

In Proposition 3.5, the 3 centers of the blowing up lie on a section of F_b if and only if the surface S' is isomorphic to S'_{I_m} for some I_m . This completes the proof of (4) and (5) of Theorem 1.5.

4. PROOF OF PROPOSITION 3.4

As before, let p_1, p_2, p_3, p_4 be the singular points of S of order 2, 3, 5, $120m$, respectively, and let $f : S' \rightarrow S$ be a minimal resolution. Let R_{p_i} be the sublattice of $H^2(S', \mathbb{Z})$ generated by all exceptional curves contained in $f^{-1}(p_i)$.

Let C be an irreducible curve on S' . By Lemma 2.5(6), C can be written as $C = kM + r$ for some integer k and some $r \in R$, hence as

$$(4.1) \quad C = \frac{k}{\sqrt{D}} f^* K_S + C(1) + C(2) + C(3) + C(4)$$

where $C(i)$ is a $\mathbb{Q}$ -divisor supported on $f^{-1}(p_i)$ that is of the form

$$C(i) = a_i e_i + r_i$$

for some integer a_i and some $r_i \in R_{p_i}$, where e_i is a generator of the discriminant group $\text{disc}(R_{p_i})$.

Lemma 4.1. *Let C be an irreducible curve on S' of the form (4.1).*

- (1) $C(i)^2 = 0$ if and only if $C(i) = 0$ if and only if C does not meet $f^{-1}(p_i)$.
- (2) $C(1)^2 = -\frac{1}{2}x$ for some integer $x \geq 0$,
 $C(1)^2 = -\frac{1}{2}$ if and only if C meets with multiplicity 1 the component of $f^{-1}(p_1)$.
- (3) Assume that p_2 is of type $< 3, 2 >$. Then
 $C(2)^2 = -\frac{2}{3}y$ for some integer $y \geq 0$,
 $C(2)^2 = -\frac{2}{3}$ if and only if C meets with multiplicity 1 exactly one of the two components of $f^{-1}(p_2)$.
- (4) Assume that p_3 is of type $< 5, 4 >$. Then
 $C(3)^2 \leq -\frac{4}{5}$ if $C(3) \neq 0$,
 $C(3)^2 = -\frac{4}{5}$ if and only if C meets with multiplicity 1 exactly one of the two end components of $f^{-1}(p_3)$.

Proof. (1) The first equivalence follows from the negative definiteness of exceptional curves. Note that $EC = EC(i)$ for any curve $E \subset f^{-1}(p_i)$.

The curve C does not meet $f^{-1}(p_i) \Leftrightarrow EC = 0$ for any curve $E \subset f^{-1}(p_i) \Leftrightarrow EC(i) = 0$ for any curve $E \subset f^{-1}(p_i) \Leftrightarrow C(i) = 0$.

(2) is trivial.

(3) Let E_1, E_2 be the exceptional curves generating R_{p_2} . Take

$$e := -\frac{E_1 + 2E_2}{3} = E_2^*$$

as a generator of $\text{disc}(R_{p_2})$. Then $C(2)$ is of the form $C(2) = ae + b_1 E_1 + b_2 E_2$ for some integers a, b_1, b_2 , hence of the form $C(2) = se + tE_2$ for some integers s, t . We

have

$$C(2)^2 = -\frac{2}{3}(s^2 - 3st + 3t^2).$$

It is easy to see that $y := s^2 - 3st + 3t^2 = (s - 3t/2)^2 + 3t^2/4 \geq 0$ for all s, t .

C meets exactly one of the two components of $f^{-1}(p_2)$ with multiplicity 1 $\Leftrightarrow (E_1C(2), E_2C(2)) = (1, 0)$ or $(0, 1) \Leftrightarrow C(2) = E_1^* = 2e + E_2$ or $C(2) = E_2^* = e \Leftrightarrow (s, t) = (2, 1)$ or $(1, 0) \Rightarrow C(2)^2 = -2/3$. Conversely, if $C(2)^2 = -2/3$, then there are six solutions $(s, t) = \pm(1, 0), \pm(2, 1), \pm(1, 1)$ for $y = (s - 3t/2)^2 + 3t^2/4 = 1$. Since $E_iC(2) = E_iC \geq 0$ for $i = 1, 2$, there remain only two solutions $(s, t) = (1, 0), (2, 1)$.

(4) Let E_1, E_2, E_3, E_4 be the exceptional curves generating R_{p_3} . Take

$$e := -\frac{E_1 + 2E_2 + 3E_3 + 4E_4}{5} = E_4^*$$

as a generator of $\text{disc}(R_{p_3})$. Then $C(3)$ is of the form $C(3) = ae + b_1E_1 + b_2E_2 + b_3E_3 + b_4E_4$ for some integers a, b_1, b_2, b_3, b_4 , hence of the form $C(3) = se + uE_2 + vE_3 + wE_4$ for some integers s, u, v, w . We have

$$\begin{aligned} C(3)^2 &= -\frac{4}{5}s^2 - 2u^2 - 2v^2 - 2w^2 + 2sw + 2uv + 2vw \\ &= -\frac{4}{5}\{(s - \frac{5w}{4})^2 + \frac{5}{2}(u - \frac{v}{2})^2 + \frac{15}{8}(v - \frac{2w}{3})^2 + \frac{5}{48}w^2\}. \end{aligned}$$

To prove the first assertion, assume that

$$(s - \frac{5w}{4})^2 + \frac{5}{2}(u - \frac{v}{2})^2 + \frac{15}{8}(v - \frac{2w}{3})^2 + \frac{5}{48}w^2 < 1.$$

We need to show that $(s, u, v, w) = (0, 0, 0, 0)$. The above inequality implies that $w^2 \leq 9$, i.e., $w = 0, \pm 1, \pm 2, \pm 3$. If $w = 0$, then there is only one solution $(s, u, v, w) = (0, 0, 0, 0)$ to the inequality. If $w = \pm 1, \pm 2, \pm 3$, no solution to the inequality. This proves the first assertion.

C meets exactly one of the two end components of $f^{-1}(p_3)$ with multiplicity 1 $\Leftrightarrow (E_1C, E_2C, E_3C, E_4C) = (1, 0, 0, 0)$ or $(0, 0, 0, 1) \Leftrightarrow C(3) = E_1^* = 4e + E_2 + 2E_3 + 3E_4$ or $C(3) = E_4^* = e \Leftrightarrow (s, u, v, w) = (4, 1, 2, 3)$ or $(1, 0, 0, 0) \Rightarrow C(3)^2 = -4/5$. Conversely, if $C(3)^2 = -4/5$, then

$$(s - \frac{5w}{4})^2 + \frac{5}{2}(u - \frac{v}{2})^2 + \frac{15}{8}(v - \frac{2w}{3})^2 + \frac{5}{48}w^2 = 1.$$

There are ten solutions

$$(s, u, v, w) = \pm(1, 0, 0, 0), \pm(4, 1, 2, 3), \pm(1, 1, 1, 1), \pm(1, 0, 1, 1), \pm(1, 0, 0, 1).$$

Since $E_iC(3) = E_iC \geq 0$ for $i = 1, 2, 3, 4$, there remain only two solutions

$$(s, u, v, w) = (4, 1, 2, 3), (1, 0, 0, 0). \quad \square$$

4.1. Case 1: $\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 4 \rangle + \langle b; 2, 1; 3, 1; 5, 1 \rangle$, $b \geq 2$. In this case, the number of exceptional curves $\mu = 11$, so $K_{S'}^2 = -2$. Let $E_1, \dots, E_4$ be the components of $f^{-1}(p_4)$ such that

$$\begin{array}{c} -3 \\ E_2 - E_4 - E_3 \\ \downarrow \\ E_1 \\ -2 \end{array}$$

We compute

$$(4.2) \quad K_{S'} = f^*K_S - \frac{(15b - 16)E_1 + (20b - 21)E_2 + (24b - 25)E_3 + (30b - 32)E_4}{30b - 31},$$

$$K_S^2 = \frac{30(b-1)^2}{30b-31}, \quad |\det(R)| = 30 \cdot (30b-31), \quad D = |\det(R)|K_S^2 = 30^2(b-1)^2.$$

We also compute the dual vectors,

$$\begin{aligned} E_1^* &= -\frac{1}{30b-31} \{(15b-8)E_1 + 5E_2 + 3E_3 + 15E_4\}, \\ E_2^* &= -\frac{1}{30b-31} \{5E_1 + (10b-7)E_2 + 2E_3 + 10E_4\}, \\ E_3^* &= -\frac{1}{30b-31} \{3E_1 + 2E_2 + (6b-5)E_3 + 6E_4\}. \end{aligned}$$

Claim 4.1.1. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then it satisfies one of the following three cases:

Case	CE_4	CE_3	CE_2	CE_1	k
(a)	0	0	0	1	-15
(b)	0	0	1	0	-10
(c)	0	1	0	0	-6

Proof. We use the same argument as in the proof of Lemma 3.1. First note that $(f^*K_S)C = \frac{k}{\sqrt{D}}(f^*K_S)^2 = \frac{(b-1)k}{30b-31}$. Since $-K_S$ is ample and $C \notin R$, $(f^*K_S)C < 0$, so $k < 0$. Intersecting C with (4.2) we get

$$C\{(15b-16)E_1 + (20b-21)E_2 + (24b-25)E_3 + (30b-32)E_4\} = (b-1)k + 30b-31.$$

This is possible only if C satisfies one of the three cases (a), (b), (c), or the case

$$(d) \quad CE_4 = 1, CE_3 = 0, CE_2 = 0, CE_1 = 0, \quad b = 2, \quad k = -1.$$

In the last case, we compute $C(4) = E_4^* = -\frac{1}{29}(15E_1 + 10E_2 + 6E_3 + 30E_4)$, so $C(4)^2 = E_4^*C(4) = -\frac{30}{29}$ and hence we get

$$\sum_{j \leq 3} C(j)^2 = C^2 - C(4)^2 - \left(\frac{-1}{30}f^*K_S\right)^2 = -1 + \frac{30}{29} - \frac{1}{30 \cdot 29} > 0,$$

contradicts the negative definiteness of exceptional curves. $\square$

Claim 4.1.2. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then C meets only one component of $f^{-1}(p_1) \cup f^{-1}(p_2) \cup f^{-1}(p_3)$, the intersection multiplicity is 1, and the component is

- (1) the component of $f^{-1}(p_1)$, if C satisfies (a),
- (2) one of the two components of $f^{-1}(p_2)$, if C satisfies (b),
- (3) one of the two end components of $f^{-1}(p_3)$, if C satisfies (c).

Proof. Assume that C satisfies (a). Then, $C(4) = E_1^*$, $C(4)^2 = E_1^*C(4) = -\frac{15b-8}{30b-31}$, $C(1)^2 + C(2)^2 + C(3)^2 = C^2 - C(4)^2 - \left(\frac{-15}{30(b-1)}f^*K_S\right)^2 = -\frac{1}{2}$.

By Lemma 4.1, $C(2) = C(3) = 0$, $C(1)^2 = -\frac{1}{2}$, and C does not meet $f^{-1}(p_2) \cup f^{-1}(p_3)$, but meets the component of $f^{-1}(p_1)$ with multiplicity 1.

Assume that C satisfies (b). Then, $C(4) = E_2^*$, $C(4)^2 = E_2^*C(4) = -\frac{10b-7}{30b-31}$, $C(1)^2 + C(2)^2 + C(3)^2 = C^2 - C(4)^2 - \left(\frac{-10}{30(b-1)}f^*K_S\right)^2 = -\frac{2}{3}$.

By Lemma 4.1, $C(1) = C(3) = 0$, $C(2)^2 = -\frac{2}{3}$, and C does not meet $f^{-1}(p_1) \cup f^{-1}(p_3)$, but meets one of the two components of $f^{-1}(p_2)$ with multiplicity 1.

Assume that C satisfies (c). Then, $C(4) = E_3^*$, $C(4)^2 = E_3^*C(4) = -\frac{6b-5}{30b-31}$, $C(1)^2 + C(2)^2 + C(3)^2 = C^2 - C(4)^2 - \left(\frac{-6}{30(b-1)}f^*K_S\right)^2 = -\frac{4}{5}$.

By Lemma 4.1, $C(1) = C(2) = 0$, $C(3)^2 = -\frac{4}{5}$, and C does not meet $f^{-1}(p_1) \cup f^{-1}(p_2)$, but meets one of the end components of $f^{-1}(p_3)$ with multiplicity 1. $\square$

Claim 4.1.3. There are three, mutually disjoint, (-1) -curves C_1, C_2, C_3 satisfying (a), (b), (c) from Claim 4.1.1, respectively.

Proof. By Lemma 3.3, S' is a rational surface. Since $K_{S'}^2 < 8$, S' contains a (-1) -curve and can be blown down to a minimal rational surface F_n or $\mathbb{CP}^2$.

Assume that there is no (-1) -curve $C \subset S'$ meeting $f^{-1}(p_4)$. Then, since S' cannot contain a $(-l)$ -curve with $l \geq 2$ other than the exceptional curves of f (Lemma 2.1), the configuration of $f^{-1}(p_4)$ remains the same under the blow down process to F_n or $\mathbb{CP}^2$. This is impossible, as the configuration would define a negative definite sublattice of rank 4 inside the Picard lattice of F_n or $\mathbb{CP}^2$.

Assume that there is only one (-1) -curve meeting $f^{-1}(p_4)$. Then, the 3 components of $f^{-1}(p_4)$ untouched by the (-1) -curve remain the same under the blow down process and define a negative definite sublattice of rank 3 inside the Picard lattice of F_n or $\mathbb{CP}^2$. This is impossible.

If there are only two (-1) -curve meeting $f^{-1}(p_4)$. Then the 2 components of $f^{-1}(p_4)$ untouched by the two (-1) -curves would remain the same under the blow down process and define a negative definite sublattice of rank 2 inside the Picard lattice of F_n or $\mathbb{CP}^2$. Again, this is impossible.

For the mutual disjointness, we note that

$$C_1 = \frac{-15}{30(b-1)} f^* K_S + C_1(1) + E_1^*,$$

$$C_2 = \frac{-10}{30(b-1)} f^* K_S + C_2(2) + E_2^*,$$

$$C_3 = \frac{-6}{30(b-1)} f^* K_S + C_3(3) + E_3^*.$$

A direct calculation shows that $C_i C_j = 0$ for $i \neq j$. □

4.2. Case 2: $\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 2 \rangle + \langle b; 2, 1; 3, 1; 5, 3 \rangle$, $b \geq 2$. In this case, $\mu = 10$, so $K_{S'}^2 = -1$. Let B_1, B_2 be the components of $f^{-1}(p_3)$, and $E_1, \dots, E_5$ be the components of $f^{-1}(p_4)$ such that

$$\begin{array}{ccccccc} -2 & -3 & -3 & -b & -2 & -3 \\ B_1 & - B_2 & E_2 & - E_5 & - E_4 & - E_3 \\ & & & \downarrow & & \\ & & & E_1 & & \\ & & & -2 & & \end{array}$$

Then

$$(4.3) \quad \begin{aligned} K_{S'} &= f^* K_S - \frac{1}{5}(B_1 + 2B_2) - \frac{1}{30b-43} \{ (15b-22)E_1 + (20b-29)E_2 \\ &\quad + (18b-26)E_3 + (24b-35)E_4 + (30b-44)E_5 \}, \end{aligned}$$

$$K_S^2 = \frac{6(5b-7)^2}{5(30b-43)}, \quad |\det(R)| = 30 \cdot (30b-43), \quad D = 6^2(5b-7)^2.$$

We also compute the dual vectors,

$$\begin{aligned} B_1^* &= -\frac{3B_1+B_2}{5} & B_2^* &= -\frac{B_1+2B_2}{5}, \\ E_1^* &= -\frac{1}{30b-43} \{ (15b-14)E_1 + 5E_2 + 3E_3 + 9E_4 + 15E_5 \}, \\ E_2^* &= -\frac{1}{30b-43} \{ 5E_1 + (10b-11)E_2 + 2E_3 + 6E_4 + 10E_5 \}, \\ E_3^* &= -\frac{1}{30b-43} \{ 3E_1 + 2E_2 + (12b-16)E_3 + (6b-5)E_4 + 6E_5 \}. \end{aligned}$$

Claim 4.2.1. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then it satisfies one of the following three cases:

Case	CE_5	CE_4	CE_3	CE_2	CE_1	CB_2	CB_1	k
(a)	0	0	0	0	1	0	0	-15
(b)	0	0	0	1	0	0	0	-10
(c)	0	0	1	0	0	0	1	-6

Proof. First note that $(f^*K_S)C = \frac{k}{\sqrt{D}}(f^*K_S)^2 = \frac{(5b-7)k}{5(30b-43)}$. Since $-K_S$ is ample and $C \notin R$, we see that $k < 0$. Intersecting C with (4.3) we get $(30b-43)C(B_1+2B_2) + 5C\{(15b-22)E_1 + (20b-29)E_2 + (18b-26)E_3 + (24b-35)E_4 + (30b-44)E_5\} = (5b-7)k + 5(30b-43) < 5(30b-43)$.

This is possible only if C satisfies one of the three cases or the following case

(d) $CE_5 = 0, CE_4 = 1, CE_3 = CE_2 = CE_1 = 0, CB_1 = 1, CB_2 = 0, b = 2, k = -1$.

In case (d), $C(3) = B_1^*$ and $C(4) = E_4^* = -\frac{1}{17}(9E_1 + 6E_2 + 7E_3 + 21E_4 + 18E_5)$, thus $C(1)^2 + C(2)^2 = C^2 - C(3)^2 - C(4)^2 - (\frac{-1}{18}f^*K_S)^2 = -1 + \frac{3}{5} + \frac{21}{17} - \frac{1}{30 \cdot 17} > 0$, contradicts the negative definiteness of exceptional curves. $\square$

Claim 4.2.2. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then C meets only one component of $f^{-1}(p_1) \cup f^{-1}(p_2) \cup f^{-1}(p_3)$, the intersection multiplicity is 1, and the component is

- (1) the component of $f^{-1}(p_1)$, if C satisfies (a),
- (2) one of the two components of $f^{-1}(p_2)$, if C satisfies (b),
- (3) the component B_1 of $f^{-1}(p_3)$, if C satisfies (c).

Proof. Assume that C satisfies (a). Then, $C(3) = 0$ and $C(4) = E_1^*$, so

$$C(1)^2 + C(2)^2 = C^2 - C(4)^2 - (\frac{-15}{6(5b-7)}f^*K_S)^2 = -\frac{1}{2}.$$

By Lemma 4.1, $C(2) = 0$ and $C(1)^2 = -\frac{1}{2}$.

Assume that C satisfies (b). Then, $C(3) = 0$ and $C(4) = E_2^*$, so

$$C(1)^2 + C(2)^2 = C^2 - C(4)^2 - (\frac{-10}{6(5b-7)}f^*K_S)^2 = -\frac{2}{3}.$$

By Lemma 4.1, $C(1) = 0$ and $C(2)^2 = -\frac{2}{3}$.

Assume that C satisfies (c). Then, $C(3) = B_1^* = -\frac{3B_1+B_2}{5}$ and $C(4) = E_3^*$, so

$$C(1)^2 + C(2)^2 = C^2 - C(3)^2 - C(4)^2 - (\frac{-6}{6(5b-7)}f^*K_S)^2 = 0.$$

By the negative definiteness, $C(1) = C(2) = 0$. $\square$

By the same proof as in the previous case, we see that there are three, mutually disjoint, (-1) -curves C_1, C_2, C_3 satisfying (a), (b), (c) from Claim 4.2.1, respectively.

4.3. Case 3: $\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 3 \rangle + \langle b, 2, 1; 3, 1; 5, 2 \rangle$, $b \geq 2$. In this case, $\mu = 10$, so $K_{S'}^2 = -1$. Let B_1, B_2 be the components of $f^{-1}(p_3)$, and $E_1, \dots, E_5$ be the components of $f^{-1}(p_4)$ such that

$$\begin{array}{ccccccc} -2 & -3 & -3 & -b & -3 & -2 \\ B_1 & -B_2 & E_2 & -E_5 & -E_4 & -E_3 \\ & & & \downarrow & & \\ & & & E_1 & & \\ & & & -2 & & \end{array}$$

Then

$$(4.4) \quad K_{S'} = f^*K_S - \frac{1}{5}(B_1 + 2B_2) - \frac{1}{30b-37}\{(15b-19)E_1 + (20b-25)E_2 + (12b-15)E_3 + (24b-30)E_4 + (30b-38)E_5\},$$

$$K_S^2 = \frac{6(5b-6)^2}{5(30b-37)}, \quad |\det(R)| = 30 \cdot (30b-37), \quad D = 6^2(5b-6)^2.$$

We also compute the dual vectors,

$$\begin{aligned} B_1^* &= -\frac{3B_1+B_2}{5}, & B_2^* &= -\frac{B_1+2B_2}{5}, \\ E_1^* &= -\frac{1}{30b-37}\{(15b-11)E_1 + 5E_2 + 3E_3 + 6E_4 + 15E_5\}, \\ E_2^* &= -\frac{1}{30b-37}\{5E_1 + (10b-9)E_2 + 2E_3 + 4E_4 + 10E_5\}, \\ E_3^* &= -\frac{1}{30b-37}\{3E_1 + 2E_2 + (18b-21)E_3 + (6b-5)E_4 + 6E_5\}. \end{aligned}$$

Claim 4.3.1. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then it satisfies one of the following three cases:

Case	CE_5	CE_4	CE_3	CE_2	CE_1	CB_2	CB_1	k
(a)	0	0	0	0	1	0	0	-15
(b)	0	0	0	1	0	0	0	-10
(c)	0	0	1	0	0	1	0	-6

Proof. Since $(f^*K_S)C = \frac{(5b-6)k}{5(30b-37)} < 0$, $k < 0$. Intersecting C with (4.4) we get $(30b-37)C(B_1+2B_2) + 5C\{(15b-19)E_1 + (20b-25)E_2 + (12b-15)E_3 + (24b-30)E_4 + (30b-38)E_5\} = (5b-6)k + 5(30b-37) < 5(30b-37)$.

This is possible only if C satisfies one of the three cases or the following case

$$(d) \quad CE_5 = 0, CE_4 = 0, CE_3 = 1, CE_2 = 0, CE_1 = 0, CB_1 = 2, CB_2 = 0, \quad k = -6.$$

In the last case, $C(3) = 2B_1^*$ and $C(4) = E_3^*$, so $C(3)^2 = -\frac{12}{5}$ and $C(4)^2 = -\frac{18b-21}{30b-37}$, hence $C(1)^2 + C(2)^2 = C^2 - C(3)^2 - C(4)^2 - (\frac{-6}{6(5b-6)}f^*K_S)^2 > 0$, which contradicts the negative definiteness of exceptional curves. $\square$

Claim 4.3.2. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then C meets only one component of $f^{-1}(p_1) \cup f^{-1}(p_2) \cup f^{-1}(p_3)$, the intersection multiplicity is 1, and the component is

- (1) the component of $f^{-1}(p_1)$, if C satisfies (a),
- (2) one of the two components of $f^{-1}(p_2)$, if C satisfies (b),
- (3) the component B_2 of $f^{-1}(p_3)$, if C satisfies (c).

Proof. Assume that C satisfies (a). Then, $C(3) = 0$ and $C(4) = E_1^*$, so $C(1)^2 + C(2)^2 = -1 + \frac{15b-11}{30b-37} - (\frac{-15}{6(5b-6)}f^*K_S)^2 = -\frac{1}{2}$. By Lemma 4.1, $C(2) = 0$ and $C(1)^2 = -\frac{1}{2}$.

Assume that C satisfies (b). Then, $C(3) = 0$ and $C(4) = E_2^*$, so $C(1)^2 + C(2)^2 = -1 + \frac{10b-9}{30b-37} - (\frac{-10}{6(5b-6)}f^*K_S)^2 = -\frac{2}{3}$. By Lemma 4.1, $C(1) = 0$ and $C(2)^2 = -\frac{2}{3}$.

Assume that C satisfies (c). Then, $C(3) = B_2^*$ and $C(4) = E_3^*$, so $C(1)^2 + C(2)^2 = -1 + \frac{2}{5} + \frac{18b-21}{30b-37} - (\frac{-6}{6(5b-6)}f^*K_S)^2 = 0$. By the negative definiteness, $C(1) = C(2) = 0$. $\square$

The same proof as in the previous cases shows that there are three, mutually disjoint, (-1) -curves C_1, C_2, C_3 satisfying (a), (b), (c) from Claim 4.3.1, respectively.

4.4. **Case 4:** $\langle 2, 1 \rangle + \langle 3, 2 \rangle + \langle 5, 1 \rangle + \langle b; 2, 1; 3, 1; 5, 4 \rangle$, $b \geq 2$. In this case, $\mu = 11$, so $K_{S'}^2 = -2$. Let B be the component of $f^{-1}(p_3)$, and $E_1, \dots, E_7$ be the components of $f^{-1}(p_4)$ such that

$$\begin{array}{c} \begin{array}{ccccccc} -3 & -b & -2 & -2 & -2 & -2 \\ E_2 & -E_7 & -E_6 & -E_5 & -E_4 & -E_3 \end{array} \\ | \\ E_1 \\ -2 \end{array}$$

Then

(4.5)

$$K_{S'} = f^*K_S - \frac{3}{5}B - \frac{1}{30b-49}\{(15b-25)E_1 + (20b-33)E_2 + (6b-10)E_3 + (12b-20)E_4 + (18b-30)E_5 + (24b-40)E_6 + (30b-50)E_7\},$$

$$K_S^2 = \frac{6(5b-8)^2}{5(30b-49)}, \quad |\det(R)| = 30 \cdot (30b-49), \quad D = 6^2(5b-8)^2.$$

We also compute the dual vectors,

$$\begin{aligned} E_1^* &= -\frac{1}{30b-49}\{(15b-17)E_1 + 5E_2 + 3E_3 + 6E_4 + 9E_5 + 12E_6 + 15E_7\}, \\ E_2^* &= -\frac{1}{30b-49}\{5E_1 + (10b-13)E_2 + 2E_3 + 4E_4 + 6E_5 + 8E_6 + 10E_7\}, \\ E_3^* &= -\frac{1}{30b-49}\{3E_1 + 2E_2 + (24b-38)E_3 + (18b-27)E_4 + (12b-16)E_5 + (6b-5)E_6 + 6E_7\}. \end{aligned}$$

Claim 4.4.1. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then it satisfies one of the following three cases:

Case	CE_7	CE_6	CE_5	CE_4	CE_3	CE_2	CE_1	CB	k
(a)	0	0	0	0	0	0	1	0	-15
(b)	0	0	0	0	0	1	0	0	-10
(c)	0	0	0	0	1	0	0	1	-6

Proof. Since $(f^*K_S)C = \frac{(5b-8)k}{5(30b-49)} < 0$, $k < 0$. Intersecting C with (4.5) we get $3(30b-49)CB + 5C\{(15b-25)E_1 + (20b-33)E_2 + (6b-10)E_3 + (12b-20)E_4 + (18b-30)E_5 + (24b-40)E_6 + (30b-50)E_7\} = (5b-8)k + 5(30b-49) < 5(30b-49)$. This is possible only if C satisfies one of the three cases, or one of the two cases:

Case	CE_7	CE_6	CE_5	CE_4	CE_3	CE_2	CE_1	CB	k	b
(d)	0	0	0	0	2	0	0	1	-1	2
(e)	0	0	0	1	0	0	0	1	-1	2

In Case (d), $C(3) = B^* = -\frac{1}{5}B$ and $C(4) = 2E_3^*$, thus

$$C(1)^2 + C(2)^2 = C^2 - C(3)^2 - C(4)^2 - (\frac{-1}{12}f^*K_S)^2 = -1 + \frac{1}{5} + \frac{40}{11} - \frac{1}{30 \cdot 11} > 0.$$

In Case (e), $C(3) = -\frac{1}{5}B$ and $C(4) = E_4^* = -\frac{1}{11}(6E_1 + 4E_2 + 9E_3 + 18E_4 + 16E_5 + 14E_6 + 12E_7)$, thus

$$C(1)^2 + C(2)^2 = C^2 - C(3)^2 - C(4)^2 - (\frac{-1}{12}f^*K_S)^2 = -1 + \frac{1}{5} + \frac{18}{11} - \frac{1}{30 \cdot 11} > 0.$$

Both contradict the negative definiteness of exceptional curves. $\square$

Claim 4.4.2. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then C meets only one component of $f^{-1}(p_1) \cup f^{-1}(p_2) \cup f^{-1}(p_3)$, the intersection multiplicity with the component is 1, and the component is

- (1) the component of $f^{-1}(p_1)$, if C satisfies (a),
- (2) one of the two components of $f^{-1}(p_2)$, if C satisfies (b),
- (3) the component B of $f^{-1}(p_3)$, if C satisfies (c).

Proof. Assume that C satisfies (a). Then, $C(3) = 0$ and $C(4) = E_1^*$, so $C(4)^2 = -\frac{15b-17}{30b-49}$ and $C(1)^2 + C(2)^2 = C^2 - C(4)^2 - (\frac{-15}{6(5b-8)}f^*K_S)^2 = -\frac{1}{2}$. By Lemma 4.1, $C(2) = 0$ and $C(1)^2 = -\frac{1}{2}$.

Assume that C satisfies (b). Then, $C(3) = 0$ and $C(4) = E_2^*$, so $C(1)^2 + C(2)^2 = -1 + \frac{10b-13}{30b-49} - (\frac{-10}{6(5b-8)}f^*K_S)^2 = -\frac{2}{3}$. By Lemma 4.1, $C(1) = 0$ and $C(2)^2 = -\frac{2}{3}$.

Assume that C satisfies (c). Then, $C(3) = -\frac{1}{5}B$ and $C(4) = E_3^*$, so $C(1)^2 + C(2)^2 = -1 + \frac{1}{5} + \frac{24b-38}{30b-49} - (\frac{-6}{6(5b-8)}f^*K_S)^2 = 0$. By the negative definiteness, $C(1) = C(2) = 0$. $\square$

The same proof as in the previous cases shows that there are three, mutually disjoint, (-1) -curves C_1, C_2, C_3 satisfying (a), (b), (c) from Claim 4.4.1, respectively.

4.5. Case 5: $\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 4 \rangle + \langle b, 2, 1; 3, 2; 5, 1 \rangle$, $b \geq 2$. In this case, $\mu = 11$, so $K_{S'}^2 = -2$. Let B be the component of $f^{-1}(p_2)$, and $E_1, \dots, E_5$ be the components of $f^{-1}(p_4)$ such that

$$\begin{array}{c} -2 \\ E_2 - E_3 - E_5 - E_4 \\ \downarrow \\ E_1 \\ -2 \end{array}$$

Then

$$(4.6) \quad K_{S'} = f^*K_S - \frac{1}{3}B - \frac{1}{30b-41}\{(15b-21)E_1 + (10b-14)E_2 + (20b-28)E_3 + (24b-33)E_4 + (30b-42)E_5\},$$

$$K_S^2 = \frac{10(3b-4)^2}{3(30b-41)}, \quad |\det(R)| = 30 \cdot (30b-41), \quad D = 10^2(3b-4)^2.$$

We also compute the dual vectors,

$$\begin{aligned} E_1^* &= -\frac{1}{30b-41}\{(15b-13)E_1 + 5E_2 + 10E_3 + 3E_4 + 15E_5\}, \\ E_2^* &= -\frac{1}{30b-41}\{5E_1 + (20b-24)E_2 + (10b-7)E_3 + 2E_4 + 10E_5\}, \\ E_4^* &= -\frac{1}{30b-41}\{3E_1 + 2E_2 + 4E_3 + (6b-7)E_4 + 6E_5\}. \end{aligned}$$

Claim 4.5.1. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then it satisfies one of the following three cases:

Case	CE_5	CE_4	CE_3	CE_2	CE_1	CB	k
(a)	0	0	0	0	1	0	-15
(b)	0	0	0	1	0	1	-10
(c)	0	1	0	0	0	0	-6

Proof. Since $(f^*K_S)C = \frac{(3b-4)k}{3(30b-41)} < 0$, $k < 0$. Intersecting C with (4.6) we get $(30b-41)CB + 3C\{(15b-21)E_1 + (10b-14)E_2 + (20b-28)E_3 + (24b-33)E_4 + (30b-42)E_5\} = (3b-4)k + 3(30b-41) < 3(30b-41)$.

This is possible only if C satisfies one of the three cases, or one of the following three cases:

Case	CE_6	CE_5	CE_4	CE_3	CE_2	CE_1	CB	k	b
(d)	0	0	0	1	0	0	1	-1	2
(e)	0	0	0	0	2	0	1	-1	2
(f)	0	0	0	0	1	1	0	-6	2

In Case (d), $C(2) = -\frac{1}{3}B$ and $C(4) = E_3^* = -\frac{1}{19}(10E_1 + 13E_2 + 26E_3 + 4E_4 + 20E_5)$, thus $C(1)^2 + C(3)^2 = C^2 - C(2)^2 - C(4)^2 - (\frac{-1}{20}f^*K_S)^2 = -1 + \frac{1}{3} + \frac{26}{19} - \frac{1}{30 \cdot 19} > 0$.

In Case (e), $C(2) = -\frac{1}{3}B$ and $C(4) = 2E_2^*$, thus

$$C(1)^2 + C(3)^2 = -1 + \frac{1}{3} + \frac{64}{19} - \frac{1}{30 \cdot 19} > 0.$$

In Case (f), $C(2) = 0$ and $C(4) = E_1^* + E_2^*$, thus $C(1)^2 + C(3)^2 = -1 + \frac{43}{19} - \frac{36}{30 \cdot 19} > 0$. All contradict the negative definiteness of exceptional curves. $\square$

Claim 4.5.2. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then C meets only one component of $f^{-1}(p_1) \cup f^{-1}(p_2) \cup f^{-1}(p_3)$, the intersection multiplicity with the component is 1, and the component is

- (1) the component of $f^{-1}(p_1)$, if C satisfies (a),
- (2) the component of B of $f^{-1}(p_2)$, if C satisfies (b),
- (3) one of the two end components of $f^{-1}(p_3)$, if C satisfies (c).

Proof. Assume that C satisfies (a). Then, $C(2) = 0$ and $C(4) = E_1^*$, so $C(4)^2 = -\frac{15b-13}{30b-41}$, hence $C(1)^2 + C(3)^2 = C^2 - C(4)^2 - (\frac{-15}{10(3b-4)}f^*K_S)^2 = -\frac{1}{2}$. By Lemma 4.1, $C(3) = 0$ and $C(1)^2 = -\frac{1}{2}$.

Assume that C satisfies (b). Then, $C(2) = -\frac{1}{3}B$ and $C(4) = E_2^*$, so $C(1)^2 + C(3)^2 = -1 + \frac{1}{3} + \frac{20b-24}{30b-41} - (\frac{-10}{10(3b-4)}f^*K_S)^2 = 0$. By the negative definiteness, $C(1) = C(3) = 0$.

Assume that C satisfies (c). Then, $C(2) = 0$ and $C(4) = E_4^*$, so $C(1)^2 + C(3)^2 = -1 + \frac{6b-7}{30b-41} - (\frac{-6}{10(3b-4)}f^*K_S)^2 = -\frac{4}{5}$. By Lemma 4.1, $C(1) = 0$ and $C(3)^2 = -\frac{4}{5}$. $\square$

The same proof as in the previous cases shows that there are three, mutually disjoint, (-1) -curves C_1, C_2, C_3 satisfying (a), (b), (c) from Claim 4.5.1, respectively. In this case, $C_1 = \frac{-15}{10(3b-4)}f^*K_S + C_1(1) + E_1^*$, $C_2 = \frac{-10}{10(3b-4)}f^*K_S + C_2(2) + E_2^*$, $C_3 = \frac{-6}{10(3b-4)}f^*K_S + C_3(3) + E_4^*$.

4.6. Case 6: $\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 2 \rangle + \langle b, 2, 1; 3, 2; 5, 3 \rangle$, $b \geq 2$. In this case, $\mu = 10$, so $K_{S'}^2 = -1$. Let B be the component of $f^{-1}(p_2)$, B_2, B_3 be the components of $f^{-1}(p_3)$, and $E_1, \dots, E_6$ be the components of $f^{-1}(p_4)$ such that

$$\begin{array}{ccccccc} -2 & -3 & -2 & -2 & -b & -2 & -3 \\ B_2 & -B_3 & E_2 & -E_3 & -E_6 & -E_5 & -E_4 \\ & & & & \downarrow & & \\ & & & & E_1 & & \\ & & & & -2 & & \end{array}$$

Then

$$\begin{aligned} (4.7) \quad K_{S'} &= f^*K_S - \frac{1}{3}B - \frac{1}{5}(B_2 + 2B_3) - \frac{1}{30b-53}\{(15b-27)E_1 + (10b-18)E_2 \\ &\quad + (20b-36)E_3 + (18b-32)E_4 + (24b-43)E_5 + (30b-54)E_6\}, \\ K_S^2 &= \frac{2(15b-26)^2}{15(30b-53)}, \quad |\det(R)| = 30 \cdot (30b-53), \quad D = 2^2(15b-26)^2. \end{aligned}$$

We also compute the dual vectors,

$$\begin{aligned} B_2^* &= -\frac{3B_2+B_3}{5} & B_3^* &= -\frac{B_2+2B_3}{5}, \\ E_1^* &= -\frac{1}{30b-53}\{(15b-19)E_1 + 5E_2 + 10E_3 + 3E_4 + 9E_5 + 15E_6\}, \\ E_2^* &= -\frac{1}{30b-53}\{5E_1 + (20b-32)E_2 + (10b-11)E_3 + 2E_4 + 6E_5 + 10E_6\}, \\ E_4^* &= -\frac{1}{30b-53}\{3E_1 + 2E_2 + 4E_3 + (12b-20)E_4 + (6b-7)E_5 + 6E_6\}. \end{aligned}$$

Claim 4.6.1. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then it satisfies one of the following three cases:

Case	CE_6	CE_5	CE_4	CE_3	CE_2	CE_1	CB_3	CB_2	CB	k
(a)	0	0	0	0	0	1	0	0	0	-15
(b)	0	0	0	0	1	0	0	0	1	-10
(c)	0	0	1	0	0	0	0	1	0	-6

Proof. Since $(f^*K_S)C = \frac{(15b-26)k}{15(30b-53)} < 0$, $k < 0$. Intersecting C with (4.7) we get $(30b-53)C(5B+3B_2+6B_3)+15C\{(15b-27)E_1+(10b-18)E_2+(20b-36)E_3+(18b-32)E_4+(24b-43)E_5+(30b-54)E_6\}=(15b-26)k+15(30b-53)$. This is possible only if C satisfies one of the three cases, or one of the following five cases:

Case	CE_6	CE_5	CE_4	CE_3	CE_2	CE_1	CB_3	CB_2	CB	k	b
(d)	0	0	0	0	1	0	0	3	0	-3	2
(e)	0	0	0	0	1	0	1	1	0	-3	2
(f)	0	0	0	0	0	1	0	1	1	-1	2
(g)	0	0	0	0	2	0	0	1	0	-6	2
(h)	0	0	0	1	0	0	0	1	0	-6	2

In Case (d), $C(2)=0, C(3)=3B_2^*$ and $C(4)=E_2^*$, thus

$$C(1)^2 = C^2 - C(3)^2 - C(4)^2 - \left(\frac{-3}{8}f^*K_S\right)^2 = -1 + \frac{27}{5} + \frac{8}{7} - \frac{9}{30.7} > 0.$$

In Case (e), $C(2)=0, C(3)=B_2^*+B_3^*=-\frac{4B_2+3B_3}{5}$ and $C(4)=E_2^*$, thus

$$C(1)^2 = C^2 - C(3)^2 - C(4)^2 - \left(\frac{-3}{8}f^*K_S\right)^2 = -1 + \frac{7}{5} + \frac{8}{7} - \frac{9}{30.7} > 0.$$

In Case (f), $C(2)=-\frac{1}{3}B, C(3)=B_2^*$ and $C(4)=E_1^*$, thus

$$C(1)^2 = C^2 - C(2)^2 - C(3)^2 - C(4)^2 - \left(\frac{-1}{8}f^*K_S\right)^2 = -1 + \frac{1}{3} + \frac{3}{5} + \frac{11}{7} - \frac{1}{30.7} > 0.$$

In Case (g), $C(2)=0, C(3)=B_2^*$ and $C(4)=2E_2^*$, thus

$$C(1)^2 = C^2 - C(3)^2 - C(4)^2 - \left(\frac{-6}{8}f^*K_S\right)^2 = -1 + \frac{3}{5} + \frac{32}{7} - \frac{36}{30.7} > 0.$$

In Case (h), $C(2)=0, C(3)=B_2^*$ and $C(4)=E_3^*=-\frac{1}{7}(10E_1+9E_2+18E_3+4E_4+12E_5+20E_6)$, thus

$$C(1)^2 = C^2 - C(3)^2 - C(4)^2 - \left(\frac{-6}{8}f^*K_S\right)^2 = -1 + \frac{3}{5} + \frac{18}{7} - \frac{36}{30.7} > 0.$$

All contradict the negative definiteness of exceptional curves. $\square$

Claim 4.6.2. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then C meets only one component of $f^{-1}(p_1) \cup f^{-1}(p_2) \cup f^{-1}(p_3)$, the intersection multiplicity with the component is 1, and the component is

- (1) the component of $f^{-1}(p_1)$, if C satisfies (a),
- (2) the component B of $f^{-1}(p_2)$, if C satisfies (b),
- (3) the component B_2 of $f^{-1}(p_3)$, if C satisfies (c).

Proof. Assume that C satisfies (a). Then, $C(2)=C(3)=0$ and $C(4)=E_1^*$, so $C(4)^2=-\frac{15b-19}{30b-53}$, hence $C(1)^2=C^2-C(4)^2-\left(\frac{-15}{2(15b-26)}f^*K_S\right)^2=-\frac{1}{2}$.

Assume that C satisfies (b). Then, $C(2)=-\frac{1}{3}B, C(3)=0$ and $C(4)=E_2^*$, so $C(1)^2=-1+\frac{1}{3}+\frac{20b-32}{30b-53}-\left(\frac{-10}{2(15b-26)}f^*K_S\right)^2=0$. Hence $C(1)=0$.

Assume that C satisfies (c). In this case, $C(2)=0, C(3)=B_2^*$ and $C(4)=E_4^*$, so $C(1)^2=-1+\frac{3}{5}+\frac{12b-20}{30b-53}-\left(\frac{-6}{2(15b-26)}f^*K_S\right)^2=0$. Hence $C(1)=0$. $\square$

The same proof as in the previous cases shows that there are three, mutually disjoint, (-1) -curves C_1, C_2, C_3 satisfying (a), (b), (c) from Claim 4.6.1, respectively.

4.7. Case 7: $\langle 2, 1 \rangle + \langle 3, 1 \rangle + \langle 5, 3 \rangle + \langle b; 2, 1; 3, 2; 5, 2 \rangle$, $b \geq 2$. In this case, $\mu = 10$, so $K_S^2 = -1$. Let B be the component of $f^{-1}(p_2)$, B_2, B_3 be the components of $f^{-1}(p_3)$, and $E_1, \dots, E_6$ be the components of $f^{-1}(p_4)$ such that

$$\begin{array}{ccccccc} -2 & -3 & -2 & -2 & -b & -3 & -2 \\ B_2 & -B_3 & E_2 & -E_3 & -E_6 & -E_5 & -E_4 \\ & & & & \downarrow & & \\ & & & & E_1 & & \\ & & & & -2 & & \end{array}$$

Then

(4.8)

$$K_{S'} = f^*K_S - \frac{1}{3}B - \frac{1}{5}(B_2 + 2B_3) - \frac{1}{30b-47}\{(15b-24)E_1 + (10b-16)E_2 + (20b-32)E_3 + (12b-19)E_4 + (24b-38)E_5 + (30b-48)E_6\},$$

$$K_S^2 = \frac{2(15b-23)^2}{15(30b-47)}, \quad |\det(R)| = 30 \cdot (30b-47), \quad D = 2^2(15b-23)^2.$$

We also compute the dual vectors,

$$\begin{aligned} B_2^* &= -\frac{3B_2+B_3}{5}, & B_3^* &= -\frac{B_2+2B_3}{5}, \\ E_1^* &= -\frac{1}{30b-47}\{(15b-16)E_1 + 5E_2 + 10E_3 + 3E_4 + 6E_5 + 15E_6\}, \\ E_2^* &= -\frac{1}{30b-47}\{5E_1 + (20b-28)E_2 + (10b-9)E_3 + 2E_4 + 4E_5 + 10E_6\}, \\ E_4^* &= -\frac{1}{30b-47}\{3E_1 + 2E_2 + 4E_3 + (18b-27)E_4 + (6b-7)E_5 + 6E_6\}. \end{aligned}$$

Claim 4.7.1. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then it satisfies one of the following three cases:

Case	CE_6	CE_5	CE_4	CE_3	CE_2	CE_1	CB_3	CB_2	CB	k
(a)	0	0	0	0	0	1	0	0	0	-15
(b)	0	0	0	0	1	0	0	0	1	-10
(c)	0	0	1	0	0	0	1	0	0	-6

Proof. Since $(f^*K_S)C = \frac{(15b-23)k}{15(30b-47)} < 0$, $k < 0$. Intersecting C with (4.8) we get $(30b-47)C(5B+3B_2+6B_3) + 15C\{(15b-24)E_1 + (10b-16)E_2 + (20b-32)E_3 + (12b-19)E_4 + (24b-38)E_5 + (30b-48)E_6\} = (15b-23)k + 15(30b-47)$.

This is possible only if C satisfies one of the three cases, or the case

(d) $CE_6 = CE_5 = 0, CE_4 = 1, CE_3 = 0, CE_2 = 1, CE_1 = 0, CB_3 = 0, CB_2 = 1, CB = 0, b = 2, k = -3$.

In the last case, $C(2) = 0, C(3) = B_2^*$ and $C(4) = E_2^* + E_4^*$, thus

$$C(1)^2 = C^2 - C(3)^2 - C(4)^2 - \left(\frac{-3}{14}f^*K_S\right)^2 = -1 + \frac{3}{5} + \frac{25}{13} - \frac{9}{30 \cdot 13} > 0,$$

which contradicts the negative definiteness of exceptional curves. $\square$

Claim 4.7.2. Let C be a (-1) -curve of the form (4.1). Suppose that C meets $f^{-1}(p_4)$. Then C meets only one component of $f^{-1}(p_1) \cup f^{-1}(p_2) \cup f^{-1}(p_3)$, the intersection multiplicity with the component is 1, and the component is

- (1) the component of $f^{-1}(p_1)$, if C satisfies (a),
- (2) the component B of $f^{-1}(p_2)$, if C satisfies (b),
- (3) the component B_3 of $f^{-1}(p_3)$, if C satisfies (c).

Proof. Assume that C satisfies (a). Then, $C(2) = C(3) = 0$ and $C(4) = E_1^*$, so $C(4)^2 = -\frac{15b-16}{30b-53}$, hence $C(1)^2 = C^2 - C(4)^2 - \left(\frac{-15}{2(15b-23)}f^*K_S\right)^2 = -\frac{1}{2}$.

Assume that C satisfies (b). Then, $C(2) = -\frac{1}{3}B, C(3) = 0$ and $C(4) = E_2^*$, so $C(1)^2 = -1 + \frac{1}{3} + \frac{20b-28}{30b-47} - \left(\frac{-10}{2(15b-23)}f^*K_S\right)^2 = 0$. Hence $C(1) = 0$.

- (1) the component of $f^{-1}(p_1)$, if C satisfies (a),

- (2) the component B of $f^{-1}(p_2)$, if C satisfies (b),
- (3) the component B_2 of $f^{-1}(p_3)$, if C satisfies (c).

Proof. Assume that C satisfies (a). Then, $C(2) = C(3) = 0$ and $C(4) = E_1^*$, so $C(4)^2 = -\frac{15b-22}{30b-59}$, hence $C(1)^2 = C^2 - C(4)^2 - (\frac{-15}{2(15b-29)}f^*K_S)^2 = -\frac{1}{2}$.

Assume that C satisfies (b). Then, $C(2) = -\frac{1}{3}B$, $C(3) = 0$ and $C(4) = E_2^*$, so $C(1)^2 = -1 + \frac{1}{3} + \frac{20b-36}{30b-59} - (\frac{-10}{2(15b-29)}f^*K_S)^2 = 0$. Hence $C(1) = 0$.

Assume that C satisfies (c). Then, $C(2) = 0$, $C(3) = -\frac{1}{5}B_2$ and $C(4) = E_4^*$, so $C(1)^2 = -1 + \frac{1}{5} + \frac{24b-46}{30b-59} - (\frac{-6}{2(15b-29)}f^*K_S)^2 = 0$. Hence $C(1) = 0$. $\square$

The same proof as in the previous cases shows that there are three, mutually disjoint, (-1) -curves C_1, C_2, C_3 satisfying (a), (b), (c) from Claim 4.8.1, respectively.

Acknowledgements. We thank János Kollár for many useful comments. We are grateful to the referee for helpful suggestions which have led to improvements in the presentation of the manuscript.

REFERENCES

- [Br] E. Brieskorn, *Rationale Singularitäten komplexer Flächen*, Invent. math. **4** (1968), 336-358.
- [FS85] R. Fintushel and R. Stern, *Pseudofree orbifolds*, Ann. of Math. (2) **122** (1985), no. 2, 335-364.
- [FS87] R. Fintushel and R. Stern, *$O(2)$ actions on the 5-sphere*, Invent. Math. **87** (1987), no. 3, 457-476.
- [HK] D. Hwang and J. Keum, *The maximum number of singular points on rational homology projective planes*, arXiv:0801.3021, to appear in J. Algebraic Geometry.
- [Keu07] J. Keum, *A rationality criterion for projective surfaces - partial solution to Kollar's conjecture*, Algebraic geometry, 75-87, Contemp. Math., **422**, Amer. Math. Soc., Providence, RI, 2007.
- [Keu08] J. Keum, *Quotients of Fake Projective Planes*, Geom. & Top. **12** (2008), 2497-2515.
- [Keu10] J. Keum, *The moduli space of $\mathbb{Q}$ -homology projective planes with 5 quotient singular points*, Acta Math. Vietnamica, **35** (2010), 79-89.
- [KM] S. Keel and J. McKernan, *Rational curves on quasi-projective surfaces*, Mem. Amer. Math. Soc. **140** (1999), no. 669.
- [KNS] R. Kobayashi, S. Nakamura, and F. Sakai *A numerical characterization of ball quotients for normal surfaces with branch loci*, Proc. Japan Acad. Ser. A, Math. Sci. **65** (1989), no. 7, 238-241.
- [Kol05] J. Kollár, *Einstein metrics on 5-dimensional Seifert bundles*, Jour. Geom. Anal. **15** (2005), no. 3, 463-495.
- [Kol08] J. Kollár, *Is there a topological Bogomolov-Miyaoka-Yau inequality?*, Pure Appl. Math. Q. (Fedor Bogomolov special issue, part I), **4** (2008), no. 2, 203-236.
- [Me] G. Megyesi, *Generalisation of the Bogomolov-Miyaoka-Yau inequality to singular surfaces*, Proc. London Math. Soc. (3) **78** (1999), 241-282.
- [Mi] Y. Miyaoka, *The maximal number of quotient singularities on surfaces with given numerical invariants*, Math. Ann. **268** (1984), 159-171.
- [MY] D. Montgomery and C. T. Yang, *Differentiable pseudo-free circle actions on homotopy seven spheres*, Proc. of the Second Conference on Compact Transformation Groups (Univ. Massachusetts, Amherst, Mass., 1971), Part I, Springer, 1972, pp. 41-101. Lecture Notes in Math., Vol. 298.
- [P] T. Petrie, *Equivariant quasi-eivalence, transversality, and normal cobordism*, Proc. Int. Cong. Math., Vancouver, 1974, pp.537-541.
- [S] F. Sakai, *Semistable curves on algebraic surfaces and logarithmic pluricanonical maps*, Math. Ann. **254** (1980), no. 2, 89-120.
- [Se] H. Seifert, *Topologie dreidimensionaler gefaserner Räume*, Acta Math. **60** (1932), 147-238.

DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, DAEJON, KOREA

Current address: School of Mathematics, Korea Institute For Advanced Study, Seoul 130-722, Korea

E-mail address: `dshwang@kias.re.kr`

SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY, SEOUL 130-722, KOREA

E-mail address: `jhkeum@kias.re.kr`